

SIMPLY CONNECTED MINIMAL SYMPLECTIC 4-MANIFOLDS WITH SIGNATURE LESS THAN -1

ANAR AKHMEDOV, SCOTT BALDRIDGE, R. İNANÇ BAYKUR, PAUL KIRK, AND B. DOUG PARK

ABSTRACT. For each pair (e, σ) of integers satisfying $2e + 3\sigma \geq 0$, $\sigma \leq -2$, and $e + \sigma \equiv 0 \pmod{4}$, with four exceptions, we construct a minimal, simply connected symplectic 4-manifold with Euler characteristic e and signature σ . We also produce simply connected, minimal symplectic 4-manifolds with signature zero (resp. signature -1) with Euler characteristic $4k$ (resp. $4k + 1$) for all $k \geq 46$ (resp. $k \geq 49$).

1. INTRODUCTION

In [6], a closed, simply connected, minimal symplectic 4-manifold with Euler characteristic 6 and signature -2 is constructed. This manifold contains a symplectic genus 2 surface with trivial normal bundle and simply connected complement and also contains two Lagrangian tori with special properties. In this article we use this manifold and apply standard constructions to fill out the part of the symplectic geography plane corresponding to signature less than -1 . Recall that Taubes proved ([35, 33, 34], also Li-Liu [19]) that minimal simply connected symplectic 4-manifolds satisfy $2e + 3\sigma \geq 0$, where e denotes the Euler characteristic and σ the signature. Moreover, every symplectic 4-manifold satisfies $e + \sigma \equiv 0 \pmod{4}$.

Our main result is the following.

Theorem A. *Let σ and e denote integers satisfying $2e + 3\sigma \geq 0$, and $e + \sigma \equiv 0 \pmod{4}$.*

If, in addition,

$$\sigma \leq -2,$$

then there exists a simply connected minimal symplectic 4-manifold with signature σ and Euler characteristic e and odd intersection form, except possibly for (σ, e) equal to $(-3, 7)$, $(-3, 11)$, $(-5, 13)$, or $(-7, 15)$.

Date: May 5, 2007.

2000 Mathematics Subject Classification. Primary 57R17; Secondary 57M05, 57R55.

Key words and phrases. Symplectic topology, Luttinger surgery, fundamental group, 4-manifold.

A. A. was partially supported by the NSF grant FRG-0244663. S. B. was partially supported by the NSF grant DMS-0507857. R. İ. B. was partially supported by the NSF Grant DMS-0305818. P. K. was partially supported by the NSF grant DMS-0604310. B.D.P. was partially supported by an NSERC discovery grant.

In terms of c_1^2 and χ_h , we construct symplectic manifolds realizing all pairs of integers satisfying $0 \leq c_1^2 \leq 8\chi_h - 2$ except $(c_1^2, \chi_h) = (5, 1), (13, 2), (11, 2)$, and $(9, 2)$.

Using Freedman's theorem [12] and Taubes's results [33, 34] this theorem can be restated by saying that there exists a minimal symplectic manifold homeomorphic but not diffeomorphic to $m\mathbb{CP}^2 \# n\overline{\mathbb{CP}}^2$ whenever $m + 2 \leq n \leq 5m + 4$ and m is odd, except possibly for $(m, n) = (1, 4), (3, 6), (3, 8)$, or $(3, 10)$. The existence of minimal symplectic 4-manifolds homeomorphic to $m\mathbb{CP}^2 \# n\overline{\mathbb{CP}}^2$ for these four pairs remains an open problem (as far as we know).

The study of the geography problem for symplectic 4-manifolds has been an area of active study in recent years. Jongil Park pioneered the problem of systematically filling in large regions of the geography plane in a series of articles [24, 25, 28] (the smooth geography also has a long history, cf. [11]). More recent articles have focused on the problem of constructing small symplectic examples; this includes the articles [27, 10, 31, 2, 4, 3, 5, 6, 9].

The methods in this article are based on inductive constructions to produce simply connected manifolds starting with a few basic models. Although there are some formal similarities between some of the fundamental group calculations carried out in this article and those in the articles [2, 5, 6, 3, 4, 9], there is an important difference between the methods used in those articles and the methods of the present article, as we now explain.

In those articles, the mechanism used to kill fundamental groups comes down to establishing precise enough control over a group presentation to conclude that all generators die. This is a subtle process which depends critically on properly identifying words in fundamental groups, since e.g. in a group a pair of elements x, y might commute but their conjugates gxg^{-1}, hyh^{-1} need not.

By contrast, the mechanism of the present paper is much softer. We use standard symplectic constructions pioneered by Gompf [14] and Luttinger [20] to kill a generator outright; subsequent generators then are killed by a simple argument. In particular, although we are explicit and careful in our fundamental group calculations in Theorem 1, Lemma 16, and elsewhere, the reader will quickly understand that our results follow as easily if one only knows the statements up to conjugacy.

To illustrate this point, in the statement of Lemma 16, the expressions for μ_6, m_6, ℓ_6 are long, but it is straightforward to see that, *up to conjugacy*, $\mu_6 = [a_1, x_2], m_6 = y_2, \ell_6 = b_1^{-1}$. This less precise information is quite sufficient to prove the results of this article.

The construction is also suitable to fill out a large region of the geography plane starting with any given symplectic 4-manifold with given characteristic numbers and containing a square zero symplectic torus. For example, Theorem 23 roughly says that given a symplectic 4-manifold X ,

one can construct a new symplectic manifold Y with the same fundamental group as X and satisfying $c_1^2(Y) = c_1^2(X) + c$ and $\chi_h(Y) = \chi_h(X) + \chi$, for any (c, χ) in the cone $0 \leq c \leq 8\chi - 2$.

Since it is known how to produce manifolds with positive signature ([30]) we apply this result to a positive signature symplectic 4-manifold and prove the following (see Theorem 24).

Theorem B. *For all integers $k \geq 45$, there exists a simply connected minimal symplectic 4-manifold $X_{2k+1, 2k+1}$ with Euler characteristic $e = 4k + 4$ and signature $\sigma = 0$.*

For all integers $k \geq 49$ there exists a simply connected minimal symplectic 4-manifold $X_{2k-1, 2k}$ with Euler characteristic $e = 4k + 1$ and signature $\sigma = -1$.

All the manifolds we produce have odd intersection forms. Hence there remain 4 minimal simply connected symplectic odd 4-manifolds of signature less than or equal to -2 , 97 minimal simply connected symplectic odd 4-manifolds of non-positive signature, and roughly 280 minimal simply connected odd symplectic 4-manifolds of signature less than or equal to 4 yet to be constructed.

We finish this introduction with a brief description of the proofs. We start with three models, the minimal symplectic 4-manifolds B, C, D . These manifolds have Euler characteristic 6, 8, and 10 and signatures $-2, -4$, and -6 respectively. Each contains a disjoint pair of homologically independent Lagrangian tori T_1 and T_2 with nullhomotopic meridians and whose complement has fundamental group $\mathbb{Z} \oplus \mathbb{Z}$. Moreover, ± 1 Luttinger surgery (see Section 2) along certain curves on one or both of these tori yields a minimal symplectic 4-manifold.

We then produce a family $B_g, g \in \mathbb{Z}$, of minimal symplectic 4-manifolds with Euler characteristic $6 + 4g$ and signature -2 by taking a symplectic sum of B with a minimal manifold constructed from Luttinger surgery on a product of surfaces. This family B_g again contains a pair of Lagrangian tori T_1, T_2 with the same properties as those in B, C, D .

Taking the symplectic sum of many copies of B, B_g, C, D (and, if needed, the elliptic surfaces $E(k)$) along their tori and performing a $+1$ Luttinger surgery on the unused Lagrangian tori yields our even signature examples. Showing that the fundamental group vanishes is simple since the fundamental groups of B, B_g, C, D and the homomorphisms induced by the inclusions of the tori are known. Usher's theorem [36] easily implies that the result is minimal. The manifolds B, B_g, C, D contain -1 surfaces disjoint from the T_i which survive to -1 surfaces in the symplectic sum and hence the result has an odd intersection form.

Producing odd signature manifolds follows the same general approach, but requires several small model manifolds with appropriate Lagrangian tori to use as seeds for the symplectic sums. The construction is not quite as clean as in the even signature case.

We construct a minimal symplectic 4-manifold $P_{5,8}$ with fundamental group \mathbb{Z} , Euler characteristic 15, and signature -3 . We also use the minimal simply connected symplectic 4-manifold

$S_{1,1}$ constructed by Gompf with Euler characteristic 23 and signature -15 . These three manifolds each contain a Lagrangian or symplectic torus appropriate for taking symplectic sums with many copies of B, B_g, C , and D , and as in the even case this produces minimal simply connected 4-manifolds of odd signature less than or equal to -5 .

The signature -3 examples are constructed by a separate argument, and a few small examples not covered by our general construction are culled from the literature (i.e. $(\sigma, e) = (-7, 11)$, $(-13, 21)$, $(-11, 19)$, $(-5, 9)$) or constructed explicitly $((\sigma, e) = (-5, 17), (-7, 19), (-9, 21))$.

The authors would like to thank R.E. Gompf for helpful comments.

2. LUTTINGER SURGERY

Given any Lagrangian torus T in a symplectic 4-manifold M , the Darboux-Weinstein theorem [21] implies that there is a parameterization of a tubular neighborhood of $T^2 \times D^2 \rightarrow \text{nb}(T) \subset M$ such that the image of $T^2 \times \{d\}$ is Lagrangian for all $d \in D^2$. Choosing any point $d \neq 0$ in D^2 gives a push off $F_d : T \rightarrow T^2 \times \{d\} \subset M - T$ called the *Lagrangian push off* or *Lagrangian framing*. Given any embedded curve $\gamma \subset T$, its image $F_d(\gamma)$ is called the *Lagrangian push off* of γ .

Any curve isotopic to $\{t\} \times \partial D^2 \subset \partial(\text{nb}(T))$ will be called a *meridian* of T and typically denoted by μ_T . In this article we will typically fix a pair of embedded curves on T intersecting transversally in one point and denote the two Lagrangian push offs by m_T and ℓ_T . The triple μ_T, m_T, ℓ_T generate $H_1(\partial(\text{nb}(T)))$. Since the 3-torus has abelian fundamental group we may choose a base point t on $\partial(\text{nb}(T))$ and unambiguously refer to $\mu_T, m_T, \ell_T \in \pi_1(\partial(\text{nb}(T)), t)$.

The push offs and meridian are used to specify coordinates for a *Luttinger surgery*. This is the process of removing a tubular neighborhood of T in M and re-gluing it so that the embedded curve representing $\mu_T m_T^p \ell_T^q$ bounds a disk for some pair of integers p, q . The resulting 4-manifold admits a symplectic structure whose symplectic form is unchanged away from a neighborhood of T ([20, 1]).

When the base point x of M is chosen off the boundary of the tubular neighborhood of T , the based loops μ_T, m_T , and ℓ_T are to be joined to x by the same path in $M - T$. These curves then define elements of $\pi_1(M - T, x)$. With p, q as above, the 4-manifold resulting from torus surgery on M has fundamental group

$$\pi_1(M - T, x) / N(\mu_T m_T^p \ell_T^q),$$

where $N(\mu_T m_T^p \ell_T^q)$ denotes the normal subgroup generated by $\mu_T m_T^p \ell_T^q$.

We will only need the cases $(p, q) = (\pm 1, 0)$ or $(0, \pm 1)$ in this article, i.e. ± 1 *Luttinger surgery* along m_T or ℓ_T .

3. THE FUNDAMENTAL GROUP OF THE COMPLEMENT OF TORI IN THE PRODUCT OF SURFACES

Let F be a genus f surface, with $f \geq 2$. Choose a base point h on F and pairs $x_i, y_i, i = 1, \dots, f$ of circles forming a symplectic basis, with x_i, y_i intersecting at $h_i \in F$. Choose paths α_i from h to h_i , so that the loops

$$\tilde{x}_i = \alpha_i x_i \alpha_i^{-1} \text{ and } \tilde{y}_i = \alpha_i y_i \alpha_i^{-1}$$

generate $\pi_1(F, h)$. Let Y_i be a circle parallel to y_i which misses α_i .

Let G a genus g surface. Choose a base point k on G , and g pairs $a_1, b_1, \dots, a_g, b_g$ of circles forming a symplectic basis, with a_i, b_i intersecting at k_i . Choose paths β_i from h to h_i , so that the loops

$$\tilde{a}_i = \beta_i a_i \beta_i^{-1} \text{ and } \tilde{b}_i = \beta_i b_i \beta_i^{-1}$$

generate $\pi_1(G, k)$. Choose parallel copies A_i of a_i and B_i of b_i which miss the paths β_i . In Figure 1 we illustrate the notation when $f = 2$ and $g = 3$.




FIGURE 1. The surface $F \times G$.

The product $F \times G$ contains the union of the two symplectic surfaces $F \times \{k\} \cup \{h\} \times G$ meeting at (h, k) . There is an identification $\pi_1(F \times G, (h, k)) = \pi_1(F, h) \times \pi_1(G, k)$ which associates the loop $\tilde{x}_i \times \{k\}$ to $(\tilde{x}_i, 1)$, $\tilde{y}_i \times \{k\}$ to $(\tilde{y}_i, 1)$, $\{h\} \times \tilde{a}_i$ to $(1, \tilde{a}_i)$ and $\{h\} \times \tilde{b}_i$ to $(1, \tilde{b}_i)$. In other words, the homomorphisms induced by the inclusions $F \times \{k\} \subset F \times G$ and $\{h\} \times G \subset F \times G$ present $\pi_1(F \times G, (h, k))$ as the product of $\pi_1(F, h)$ and $\pi_1(G, k)$.

When there is no chance of confusion we denote the $2f + 2g$ loops $\tilde{x}_i \times \{k\}, \tilde{y}_i \times \{k\}, \{h\} \times \tilde{a}_i, \{h\} \times \tilde{b}_i$ simply by $\tilde{x}_i, \tilde{y}_i, \tilde{a}_i, \tilde{b}_i$. These are loops in $F \times G$ based at (h, k) .

The product $F \times G$ contains $2g$ Lagrangian tori

$$Y_1 \times A_j, \quad Y_2 \times B_j, \quad j = 1, \dots, g.$$

These $2g$ tori are pairwise disjoint and miss $(F \times \{k\}) \cup (\{h\} \times G)$.

Let N denote a tubular neighborhood of the union of these $2g$ tori:

$$N = nbd((\cup_j Y_1 \times A_j) \cup (\cup_j Y_2 \times B_j)) \subset F \times G.$$

The loops $\tilde{x}_i, \tilde{y}_i, \tilde{a}_i, \tilde{b}_i$ are loops in $F \times G - N$ based at (h, k) .

Typically, removing a surface from a 4-manifold increases the number of generators of the fundamental group, but since these tori respect the product structure one can prove the following theorem.

Theorem 1. *The $2f + 2g$ loops $\tilde{x}_1, \tilde{y}_1, \dots, \tilde{x}_f, \tilde{y}_f, \tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_g, \tilde{b}_g$ generate $\pi_1(F \times G - N, (h, k))$.*

Moreover, there are paths $d_j : [0, 1] \rightarrow F \times G - N$ from (h, k) to the boundary of the tubular neighborhood of $Y_1 \times A_j$ and $e_j : [0, 1] \rightarrow F \times G - N$ from (h, k) to the boundary of the tubular neighborhood of $Y_2 \times B_j$ so that with respect to these paths, the meridian and two Lagrangian push offs of $Y_1 \times A_j$ are homotopic in $F \times G - N$ rel endpoint to

$$\mu_{Y_1 \times A_j} = [\tilde{x}_1, \tilde{b}_j], \quad m_{Y_1 \times A_j} = \tilde{y}_1, \quad \ell_{Y_1 \times A_j} = \tilde{a}_j$$

and the meridian and two Lagrangian push offs of $Y_2 \times B_j$ are homotopic in $F \times G - N$ rel endpoint to

$$\mu_{Y_2 \times B_j} = [\tilde{x}_2, \tilde{a}_j], \quad m_{Y_2 \times B_j} = \tilde{y}_2, \quad \ell_{Y_2 \times B_j} = \tilde{b}_j.$$

Proof. Before we start the proof, we give an indication of how it will proceed. Note that $\cup_j (Y_1 \times A_j) = Y_1 \times (\cup_j A_j)$ lies on $Y_1 \times G$ and that $Y_2 \times (\cup_j B_j)$ lies on $Y_2 \times G$. Thus $F \times G - N$ can be constructed by cutting $F \times G$ along the hypersurface $(Y_1 \cup Y_2) \times G$, and then regluing the two copies of $Y_1 \times G$ only along the complement of a neighborhood of the A_i and regluing the two copies of $Y_2 \times G$ only along the complement of a neighborhood of the B_i . However, in order to use the Seifert-Van Kampen theorem, the subsets and their intersection in a decomposition are required to be connected, and so we need to modify the decomposition slightly.

Let P_1 be the annulus in F bounded by y_1 and Y_1 . Similarly let P_2 denote the annulus in F bounded by y_2 and Y_2 . Let α denote the arc $(\alpha_1 \cup \alpha_2) \times \{k\}$. Let γ_1 denote the arc $(x_1 \cap P_1) \times \{k\}$; it spans the two circles y_1 and Y_1 . Similarly let γ_2 denote the arc $(x_2 \cap P_2) \times \{k\}$. See Figure 2.




FIGURE 2.

Set

$$S_1 = (P_1 \times G) \cup \alpha \cup (P_2 \times G)$$

and

$$S_2 = (\gamma_1 \cup \gamma_2) \cup ((F - \text{int}(P_1 \cup P_2)) \times G).$$

Then in $F \times G$, $S_1 \cap S_2$ is the union of four copies of $S^1 \times G$ together with three arcs which connect the four components. In particular, S_1, S_2 and $S_1 \cap S_2$ are connected and contain the base point (h, k) .

Let $G_A = G - \text{nb}d(\cup_i A_i)$ denote the complement of an open tubular neighborhood of the A_i in G . Since the A_i do not disconnect G , G_A is path connected. Similarly let $G_B = G - \text{nb}d(\cup_i B_i)$ denote the complement of an open tubular neighborhood of the B_i in G .

To construct $F \times G - N$ we form the identification space

$$F \times G - N = S_1 \sqcup S_2 / \sim$$

by identifying $(f, s) \in S_1$ with its corresponding point (f', s') in S_2 *except if* $f \in Y_1$ and $s \in \text{nb}d(A_i)$ or $f \in Y_2$ and $s \in \text{nb}d(B_i)$. In other words, along $Y_1 \times G$ we identify only the two copies of $Y_1 \times G_A$ and along $Y_2 \times G$ we identify only the two copies of $Y_2 \times G_B$.

Hence we have exhibited $F \times G - N$ as the union of S_1 and S_2 with connected intersection

$$S_1 \cap S_2 = (Y_1 \times G_A) \cup \gamma_1 \cup (y_1 \times G) \cup \alpha \cup (y_2 \times G) \cup \gamma_2 \cup (Y_2 \times G_B).$$

It is easy to see that $\pi_1(S_1 \cap S_2, (h, k)) \rightarrow \pi_1(S_1, (h, k))$ is surjective. Indeed, one can use the product parameter in the annuli P_1 and P_2 to define a deformation retract (fixing α and hence also (h, k)) of S_1 to the subset $(y_1 \times G) \cup \alpha \cup (y_2 \times G)$ of $S_1 \cap S_2$.

The Seifert-Van Kampen theorem applies and implies that there is a surjection

$$\pi_1(S_2, (h, k)) \rightarrow \pi_1(F \times G - N, (h, k))$$

induced by inclusion.

We will show that the image of $\pi_1(S_2, (h, k)) \rightarrow \pi_1(F \times G - N, (h, k))$ is generated by the loops $\tilde{x}_1, \tilde{y}_1, \tilde{x}_2, \tilde{y}_2, \dots, \tilde{x}_f, \tilde{y}_f, \tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_g, \tilde{b}_g$. Notice that all these loops are contained in S_2 .

We find generators for $\pi_1(S_2, (h, k))$. This is again a straightforward application of the Seifert-Van Kampen theorem, as we will now show.

Since the arcs γ_i are just segments that lie on \tilde{x}_i (and the rest of the loops \tilde{x}_i lie in S_2), we can decompose S_2 as

$$S_2 = (\tilde{x}_1 \cup \tilde{x}_2) \cup ((F - \text{int}(P_1 \cup P_2)) \times G).$$

The intersection of the two pieces in this decomposition is the (contractible) set $\tilde{x}_1 \cup \tilde{x}_2 - (\gamma_1 \cup \gamma_2)$.

Hence $\pi_1(S_2, (h, k))$ is generated by \tilde{x}_1, \tilde{x}_2 and any set of generators of

$$\pi_1((F - \text{int}(P_1 \cup P_2)) \times G, (h, k)) = \pi_1(F - \text{int}(P_1 \cup P_2), h) \times \pi_1(G, k).$$

The loops \tilde{a}_i, \tilde{b}_i generate $\pi_1(G)$. The space $F - \text{int}(P_1 \cup P_2)$ is a 4-punctured genus $f - 2$ surface. Its fundamental group is generated by $\tilde{y}_1, \tilde{y}_2, \tilde{x}_3, \tilde{y}_3, \dots, \tilde{x}_f, \tilde{y}_f$ and one other loop τ based at h which is obtained by traveling from the base point to a point on the boundary component Y_1 , following Y_1 , then returning to the base point.

We have shown that the loops $\tilde{x}_1, \tilde{y}_1, \tilde{x}_2, \tilde{y}_2, \tilde{x}_3, \tilde{y}_3, \dots, \tilde{x}_f, \tilde{y}_f, \tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_g, \tilde{b}_g$, and $\tau \times \{k\}$ generate $\pi_1(S_2, (h, k))$. Hence, considered as loops in $F \times G - N$, they generate $\pi_1(F \times G - N, (h, k))$. We need only show that the generator $\tau \times \{k\}$ is not needed. But this is obvious since $\tilde{x}_1, \tilde{y}_1, \dots, \tilde{x}_f, \tilde{y}_f$ and $\tau \times \{k\}$ all lie on the surface $F \times \{k\} \subset F \times G - N$, and $\tilde{x}_1, \tilde{y}_1, \dots, \tilde{x}_f, \tilde{y}_f$ generate $\pi_1(F \times \{k\}, k)$.

We next turn to the problem of expressing the meridians and Lagrangian push offs of the generators of the Lagrangian tori $Y_1 \times A_j, Y_2 \times B_j$ in terms of the loops $\tilde{x}_1, \tilde{y}_1, \dots, \tilde{x}_f, \tilde{y}_f, \tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_g, \tilde{b}_g$. We do this for $Y_1 \times A_1$. Symmetric arguments provide the analogous calculations for the rest.

In Figure 1, denote by h_1 the intersection of x_1 and y_1 (i.e. the endpoint of α_1) and denote by k_1 the intersection of a_1 and b_1 . Then the point (h_1, k_1) lies on the boundary of the tubular neighborhood of $Y_1 \times A_1$.

Since we take the product symplectic form on $F \times G$, referring to Figure 1 one sees that the loops $y_1 \times \{k_1\}$ and $\{h_1\} \times a_1$ are Lagrangian push offs of two generators of $\pi_1(Y_1 \times A_1)$ to the boundary of the tubular neighborhood of $Y_1 \times A_1$.

There is a map of a square into $F \times G - N$ given by $\alpha_1 \times \beta_1$:

$$\alpha_1 \times \beta_1 : [0, 1] \times [0, 1] \rightarrow F \times G - N.$$

The point $(0, 0)$ is mapped to the base point (h, k) of $F \times G - N$, and the point $(1, 1)$ is mapped to (h_1, k_1) . Thus the diagonal path $d(t) = (\alpha_1(t), \beta_1(t))$ connects the base point to the boundary of the tubular neighborhood of T_1 .

Conjugating by d expresses the Lagrangian push offs as based curves in $F \times G - N$. So

$$m_{Y_1 \times A_1} = d * (y_1 \times \{k_1\}) * d^{-1} \text{ and } \ell_{Y_1 \times A_1} = d * (\{h_1\} \times a_1) * d^{-1}.$$

But $m_{Y_1 \times A_1}$ is homotopic rel basepoint in $F \times G - N$ to \tilde{y}_1 . An explicit homotopy is given by the formula:

$$(s, t) \mapsto \begin{cases} (\alpha_1(3t), \beta_1((1-s)3t)) & \text{if } 0 \leq t \leq \frac{1}{3} \\ (y_1(3t-1), \beta_1(1-s)) & \text{if } \frac{1}{3} \leq t \leq \frac{2}{3} \\ (\alpha_1(3-3t), \beta_1((1-s)(3-3t))) & \text{if } \frac{2}{3} \leq t \leq 1. \end{cases}$$

A similar homotopy, but exchanging the roles of α_1 and β_1 , establishes that $\ell_{Y_1 \times A_1}$ is homotopic rel basepoint in $F \times G - N$ to \tilde{a}_1 . (These homotopies clearly miss all the other $Y_1 \times A_j$ and $Y_2 \times B_j$.)

It remains to calculate the meridian of $Y_1 \times A_1$. For this, consider the map $x_1 \times b_1 : [0, 1] \times [0, 1] \rightarrow F \times G$. This has image a torus intersecting T_1 transversally in one point (near the point $(x_1(.9), b_1(.9))$, as one sees from Figure 1). Since

$$(h_1, k_1) = (x_1 \times b_1)(0, 0) = (x_1 \times b_1)(0, 1) = (x_1 \times b_1)(1, 0) = (x_1 \times b_1)(1, 1),$$

by conjugating the path that follows the boundary of this square by the path d from the base point (h, k) to (h_1, k_1) , we see that the meridian $\mu_{Y_1 \times A_1}$ is given by the composite

$$\mu_{Y_1 \times A_1} = d * (x_1 \times \{k_1\}) * (\{h_1\} \times b_1) * (x_1 \times \{k_1\})^{-1} * (\{h_1\} \times b_1)^{-1} * d^{-1}.$$

Now $d * (x_1 \times \{k_1\}) * d^{-1}$ is homotopic rel basepoint to \tilde{x}_1 in $F \times G - N$ by the same argument given above. The key observation is that β_1 misses A_i and B_i for all i . Similarly $d * (\{h_1\} \times b_1) * d^{-1}$ is homotopic rel basepoint to \tilde{b}_1 in $F \times G - N$. Thus

$$\mu_{Y_1 \times A_1} \sim \tilde{x}_1 * \tilde{b}_1 * \tilde{x}_1^{-1} * \tilde{b}_1^{-1} = [\tilde{x}_1, \tilde{b}_1].$$

Similar calculations establish all other assertions. \square

4. TELESCOPING TRIPLES AND SYMPLECTIC SUMS

Our construction of symplectic 4-manifolds which fill large regions in the geography plane is based on using telescoping symplectic sums along symplectic tori as well as Luttinger surgeries. The basic models in our construction have a convenient property preserved under appropriate symplectic sum, and so we formalize the property in the following definition.

Definition 2. An ordered triple (X, T_1, T_2) where X is a symplectic 4-manifold and T_1, T_2 are disjointly embedded Lagrangian tori is called a *telescoping triple* if

- (1) The tori T_1, T_2 span a 2-dimensional subspace of $H_2(X; \mathbb{R})$.
- (2) $\pi_1(X) \cong \mathbb{Z}^2$ and the inclusion induces an isomorphism $\pi_1(X - (T_1 \cup T_2)) \rightarrow \pi_1(X)$ (in particular the meridians of the T_i are trivial in $\pi_1(X - (T_1 \cup T_2))$).
- (3) The image of the homomorphism induced by inclusion $\pi_1(T_1) \rightarrow \pi_1(X)$ is a *summand* $\mathbb{Z} \subset \pi_1(X)$.
- (4) The homomorphism induced by inclusion $\pi_1(T_2) \rightarrow \pi_1(X)$ is an isomorphism.

If X is minimal we call (X, T_1, T_2) a *minimal telescoping triple*.

Note that the order of (T_1, T_2) , matters in this definition. Notice also that since the meridians $\mu_{T_1}, \mu_{T_2} \in \pi_1(X - (T_1 \cup T_2))$ are trivial and the relevant fundamental groups are abelian, the push off of an oriented loop $\gamma \subset T_i$ into $X - (T_1 \cup T_2)$ with respect to any framing of the normal bundle of T_i (e.g. the Lagrangian framing) represents a well defined element of $\pi_1(X - (T_1 \cup T_2))$, independent of the choice of framing (and basing).

The definition of a telescoping triple includes the hypothesis that the Lagrangian tori T_1 and T_2 are linearly independent in $H_2(X; \mathbb{R})$. This implies ([14]) that the symplectic form on X can be slightly perturbed so that one of the T_i remains Lagrangian and the other becomes symplectic. It can also be perturbed so that both become symplectic. Moreover, if F is a symplectic surface in X disjoint from T_1 and T_2 , the perturbed symplectic form can be chosen so that F remains symplectic.

Recall that the symplectic sum ([14]) of two symplectic 4-manifolds X and X' along genus g symplectic surfaces $F \subset X$ and $F' \subset X'$ of opposite square is a symplectic 4-manifold described topologically as the union

$$X \#_{F, F'} X' = (X - \text{nb}(F)) \cup (X' - \text{nb}(F'))$$

where the boundaries of the tubular neighborhood are identified by a fiber-preserving diffeomorphism of the corresponding circle bundles. When the surfaces are clear from context we write $X \#_s X'$.

Proposition 3. *Let (X, T_1, T_2) and (X', T'_1, T'_2) be two telescoping triples. Then for an appropriate gluing map the triple*

$$(X \#_{T_2, T'_1} X', T_1, T'_2)$$

is again a telescoping triple.

The Euler characteristic and signature of $X \#_{T_2, T'_1} X'$ are given by $e(X) + e(X')$ and $\sigma(X) + \sigma(X')$.

Proof. Let $i_j : \pi_1(T_j) \rightarrow \pi_1(X)$ be the homomorphisms induced by inclusion for $j = 1, 2$. Choose $x_1, y_1 \in \pi_1(T_1)$ so that x_1 spans the kernel of i_1 and $i_1(y_1)$ spans the image of i_1 . Denote $i_1(y_1)$ by t and choose $s \in \pi_1(X)$ so that s, t forms a basis of $\pi_1(X)$. Then choose generators x_2, y_2 for $\pi_1(T_2)$ so that $i_2(x_2) = s$ and $i_2(y_2) = t$. Thus the inclusions induce

$$x_1 \mapsto 1, y_1 \mapsto t, x_2 \mapsto s, y_2 \mapsto t.$$

Similarly, construct generators x'_1, y'_1 for $\pi_1(T'_1)$, x'_2, y'_2 for $\pi_1(T'_2)$ and s', t' for $\pi_1(X')$.

The inclusion induces an isomorphism $\pi_1(X - (T_1 \cup T_2)) \rightarrow \pi_1(X)$ and the boundary of the tubular neighborhood of T_1 (resp. T_2) is a 3-torus whose fundamental group is spanned by x_1, y_1, μ_{T_1} (resp. x_2, y_2, μ_{T_2}) (for definitiveness use the Lagrangian framing to push the x_i, y_i into the boundary of the tubular neighborhood). Similar assertions hold for (X', T'_1, T'_2) . The symplectic sum of X and X' along the surfaces $T_2 \subset X, T'_1 \subset X'$ can be formed so that the ordered triple (x_2, y_2, μ_2) is sent to (x'_1, y'_1, μ'_1) by the gluing diffeomorphism (perhaps after a change of orientation on some of the loops to ensure that the gluing diffeomorphism is orientation preserving).

The Seifert-Van Kampen theorem and the fact that all meridians are trivial imply that

$$\pi_1(X \#_{T_2, T'_1} X') = \langle s, t, s', t' \mid [s, t], [s', t'], s, t(t')^{-1} \rangle = \mathbb{Z}s' \oplus \mathbb{Z}t'.$$

The inclusion $T_1 \subset X \#_{T_2, T'_1} X'$ induces $x_1 \mapsto 1, y_1 \mapsto t'$. The inclusion $T'_2 \subset X \#_{T_2, T'_1} X'$ induces $x'_2 \mapsto s', y'_2 \mapsto t'$. Hence $(X \#_{T_2, T'_1} X', T_1, T'_2)$ is indeed a telescoping triple.

The assertions about the Euler characteristic and signature are clear. \square

Since the meridians of the Lagrangian tori are trivial in a telescoping triple, one immediately concludes the following.

Proposition 4. *Let (X, T_1, T_2) be a telescoping triple. Let ℓ_{T_1} be a Lagrangian push off of a curve on T_1 and m_{T_2} the Lagrangian push off of a curve on T_2 so that ℓ_{T_1} and m_{T_2} generate $\pi_1(X)$.*

Then the symplectic 4-manifold obtained by performing +1 Luttinger surgery on T_1 along ℓ_{T_1} and +1 surgery on T_2 along m_{T_2} is simply connected. \square

We will have frequent use of the following two results. The first is a criterion given by Usher [36] to determine when a symplectic 4-manifold is minimal. The second is a useful result of T. J. Li which we will use to verify that the hypotheses in Usher's theorem hold in certain contexts.

Theorem 5 (Usher). *Let $Z = X_1 \#_{F_1=F_2} X_2$ denote the symplectic sum of X_1 and X_2 along symplectic surfaces F_i of positive genus g . Then:*

- (i) *If either $X_1 - F_1$ or $X_2 - F_2$ contains an embedded symplectic sphere of square -1 , then Z is not minimal.*
- (ii) *If one of the summands X_i (for definiteness, say X_1) admits the structure of an S^2 -bundle over a surface of genus g such that F_1 is a section of this fiber bundle, then Z is minimal if and only if X_2 is minimal.*
- (iii) *In all other cases, Z is minimal.*

Corollary 3 of T.J. Li's article [18] provides a useful method to eliminate the first two cases of Usher's theorem in some contexts.

Theorem 6 (Li). *Let M be a symplectic 4-manifold which is not rational or ruled. Then every smoothly embedded -1 sphere is homologous to a symplectic -1 curve up to sign. If M is the blow up of a minimal symplectic 4-manifold with E_1, \dots, E_n represented by exceptional curves, then the E_i are the only classes represented by a smoothly embedded -1 sphere, hence any orientation preserving diffeomorphism maps E_i to some $\pm E_j$.*

5. THE MODEL EVEN SIGNATURE MANIFOLDS

We will setup an inductive argument by constructing telescoping symplectic sums starting with several basic telescoping triples. Proposition 4 then applies to produce simply connected 4-manifolds.

To begin with, in [6, Theorem 20], a minimal telescoping triple (B, T_1, T_2) is constructed (B is denoted B_1 in that article) so that B contains a genus 2 surface F with trivial normal bundle, and a geometrically dual symplectic -1 torus H_1 . The tori T_1, T_2 miss $F \cup H_1$. Moreover, $(B - F, T_1, T_2)$ is also a telescoping triple. These facts follow immediately from the following theorem, which summarizes the assertions established in [6].

Theorem 7. *There exists a minimal symplectic 4-manifold B containing a pair of homologically essential Lagrangian tori T_1 and T_2 and a square zero symplectic genus 2 surface F so that T_1, T_2 , and F are pairwise disjoint, $e(B) = 6$ and $\sigma(B) = -2$, and*

- (1) $\pi_1(B - (F \cup T_1 \cup T_2)) = \mathbb{Z}^2$, generated by t_1 and t_2 .
- (2) *The inclusion $B - (F \cup T_1 \cup T_2) \subset B$ induces an isomorphism on fundamental groups. In particular the meridians $\mu_F, \mu_{T_1}, \mu_{T_2}$ all vanish in $\pi_1(B - (F \cup T_1 \cup T_2))$.*
- (3) *The Lagrangian push offs m_{T_1}, ℓ_{T_1} of $\pi_1(T_1)$ are sent to 1 and t_2 respectively in the fundamental group of $B - (F \cup T_1 \cup T_2)$.*
- (4) *The Lagrangian push offs m_{T_2}, ℓ_{T_2} of $\pi_1(T_2)$ are sent to t_1 and t_2 respectively in the fundamental group of $B - (F \cup T_1 \cup T_2)$.*
- (5) *The push off $F \subset B - (F \cup T_1 \cup T_2)$ takes the first three generators of a standard symplectic generating set $\{a_1, b_1, a_2, b_2\}$ for $\pi_1(F)$ to 1 and the last element to t_2 .*
- (6) *There exists a symplectic torus $H_1 \subset B$ which intersects F transversally once, which has square -1 , and the homomorphism $\pi_1(H_1) \rightarrow \pi_1(B)$ takes the first generator to 1 and the second to t_1 . Moreover H_1 is disjoint from T_1 and T_2 (see [6, Proposition 12, Theorem 20]).*

□

The following is a restatement of [6, Theorem 13]. We state it formally since we will have frequent need of it.

Corollary 8. *The symplectic 4-manifold $X_{1,3}$ obtained from B by $+1$ Luttinger surgery on T_1 along ℓ_{T_1} and $+1$ Luttinger surgery on T_2 along m_{T_2} is a minimal symplectic 4-manifold homeomorphic to $\mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$. It contains a genus 2 symplectic surface of square zero with simply connected complement and a symplectic torus H_1 of square -1 intersecting F transversally and positively in one point.*

□

Corollary 9. *For each $g \geq 0$ there exists a minimal telescoping triple (B_g, T_1, T_2) satisfying $e(B_g) = 6 + 4g$ and $\sigma(B_g) = -2$ and containing a square -1 genus $g + 1$ surface disjoint from $T_1 \cup T_2$.*

Proof. To avoid confusing notation, during this proof we denote the symplectic genus 2 surface in B of Theorem 7 by F_B .

Take the product $F \times G$ of a genus 2 surface F and a genus g surface G , as in Section 3. Let Z_g denote the 4-manifold obtained from $F \times G$ by performing $-1/1$ Luttinger surgeries on the $2g$ disjoint Lagrangian tori $Y_1 \times A_i$ and $Y_2 \times B_i$ along the curves $\ell_{Y_1 \times A_i} = a_i$ and $\ell_{Y_2 \times B_i} = b_i$. Then by Theorem 1 the fundamental group of Z_g is generated by loops $\tilde{x}_1, \tilde{y}_1, \tilde{x}_2, \tilde{y}_2, \tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_g, \tilde{b}_g$ and the relations

$$[\tilde{x}_1, \tilde{b}_i] = \tilde{a}_i, [\tilde{x}_2, \tilde{a}_i] = \tilde{b}_i$$

hold in $\pi_1(Z_g)$. Moreover, the standard symplectic generators for $\pi_1(F)$ are sent to $\tilde{x}_1, \tilde{y}_1, \tilde{x}_2, \tilde{y}_2$ in $\pi_1(Z_g)$.

Since the meridian μ_{F_B} of $F_B \subset B$ is trivial, the symplectic sum of B with Z_g along their genus 2 symplectic surfaces $F_B \subset B$ and $F = F \times \{k\} \subset F \times G$

$$B_g = B \#_{F_B, F} Z_g$$

has fundamental group a quotient of $(\mathbb{Z}t_1 \oplus \mathbb{Z}t_2) * \pi_1(Z_g)$. We choose this symplectic sum so that the generators a_1, b_1, a_2, b_2 for $\pi_1(F_B)$ are identified (in order) with the generators $\tilde{x}_1, \tilde{y}_1, \tilde{x}_2, \tilde{y}_2$.

The fifth assertion of Theorem 7 shows that \tilde{x}_1, \tilde{y}_1 , and \tilde{x}_2 are trivial in $\pi_1(B_g)$. The relations coming from the Luttinger surgeries then show that $\tilde{a}_i = 1 = \tilde{b}_i$. Since $b_2 = \tilde{y}_2$ is identified with t_2 , $\pi_1(B_g)$ is generated by t_1 and t_2 . A calculation using the Mayer-Vietoris sequence shows that $H_1(B_g) = \mathbb{Z}^2$, and so $\pi_1(B_g) = \mathbb{Z}t_1 \oplus \mathbb{Z}t_2$. Hence (B_g, T_1, T_2) is a telescoping triple, as desired.

The Euler characteristic of B_g is calculated as

$$e(B_g) = e(B) + e(F \times G) + 4 = 6 + 4g - 4 + 4 = 6 + 4g,$$

and the signature is computed by Novikov additivity: $\sigma(B_g) = \sigma(B) = -2$.

The torus H_1 in $B - N$ geometrically dual to F_B can be lined up with one of the parallel copies $\{z\} \times G$ in $F \times G - N$ (i.e. take a *relative symplectic sum*, [14]) to produce a square -1 genus $g + 1$ surface in B_g .

Minimality follows from [6, Lemma 2], which shows that Z_g is minimal (its universal cover is contractible, so $\pi_2(Z_g) = 0$) and Usher's theorem, Theorem 5. \square

We can also produce telescoping triples with odd signature starting with B . Recall that a symplectic 4-manifold X containing a symplectic surface F is called *relatively minimal* if every -1 sphere in X intersects F .

Lemma 10. *The blow up $A = B \# \overline{\mathbb{CP}}^2$ contains a genus 3 symplectic surface F_3 with trivial normal bundle and two Lagrangian tori T_1 and T_2 so that the surfaces F_3, T_1, T_2 are pairwise disjoint, (A, F_3) is relatively minimal, and:*

- (1) $\pi_1(A - (F_3 \cup T_1 \cup T_2)) = \mathbb{Z}^2$, generated by t_1 and t_2 .
- (2) *The inclusion $A - (F_3 \cup T_1 \cup T_2) \subset A$ induces an isomorphism on fundamental groups. In particular the meridians $\mu_{F_3}, \mu_{T_1}, \mu_{T_2}$ all vanish in $\pi_1(A - (F_3 \cup T_1 \cup T_2))$.*
- (3) *The Lagrangian push offs m_{T_1}, ℓ_{T_1} of $\pi_1(T_1)$ are sent to 1 and t_2 respectively in the fundamental group of $A - (F_3 \cup T_1 \cup T_2)$.*
- (4) *The Lagrangian push offs m_{T_2}, ℓ_{T_2} of $\pi_1(T_2)$ are sent to t_1 and t_2 respectively in the fundamental group of $A - (F_3 \cup T_1 \cup T_2)$.*
- (5) *There is a standard symplectic generating set $\{a_1, b_1, a_2, b_2, a_3, b_3\}$ for $\pi_1(F_3)$ so that the push off $F_3 \subset A - (F_3 \cup T_1 \cup T_2)$ takes b_2 to t_2 , b_3 to t_1 , and all other generators to 1.*

In particular, (A, T_1, T_2) is a telescoping triple.

Proof. The 4-manifold B of Theorem 7 contains a symplectic genus 2 surface F of square zero and a geometrically dual symplectic torus H_1 of square -1 . Symplectically resolve the union $F \cup H_1$ to get F'_3 , a genus three symplectic surface in B which misses T_1 and T_2 . The surface F'_3 has square $(F + H_2)^2 = 1$. Blow up B at one point on F'_3 to construct A and denote the proper transform of F'_3 by F_3 .

Since F_3 has a geometrically dual 2-sphere (the exceptional sphere), the meridian of F_3 in $A - F_3 \subset F_3$ is nullhomotopic. The rest of the fundamental group assertions follow from Theorem 7.

Although A is not minimal, T.J. Li's theorem (Theorem 6) implies that every -1 sphere in A intersects F_3 , since B is minimal, and neither rational nor ruled. □

Note that Luttinger surgery on T_1 and T_2 in A produces a symplectic 4-manifold homeomorphic to $\mathbb{CP}^2 \# 4\overline{\mathbb{CP}}^2$, but this manifold is not minimal; it is just the blow up $X_{1,3} \# \overline{\mathbb{CP}}^2$. We do not know how to produce a minimal symplectic 4-manifold with this homeomorphism type.

We next produce a 4-manifold C with $e = 8$ and $\sigma = -2$ by stopping the construction of a minimal symplectic 4-manifold homeomorphic to $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$ in the proof of [6, Theorem 10] before the last 2 Luttinger surgeries to obtain the following.

Theorem 11. *There exists a minimal telescoping triple (C, T_1, T_2) with $e(C) = 8$ and $\text{sign}(C) = -4$. Moreover, C contains a square -1 torus disjoint from $T_1 \cup T_2$.*

Proof. We follow the notation and proof of [6, Theorem 10]. By *not* performing the Luttinger surgeries on the tori T_3 and T_4 , one obtains a minimal symplectic 4-manifold C such that $\pi_1(C - (T_3 \cup T_4))$ is generated by the two commuting elements y and a_2 . The Mayer-Vietoris sequence shows that $H_1(C - (T_3 \cup T_4); \mathbb{Z}) = \mathbb{Z}^2$, and so $\pi_1(C - (T_1 \cup T_2)) = \mathbb{Z}y \oplus \mathbb{Z}a_2$. The meridians and Lagrangian push offs of T_3 and T_4 are given by $\mu_{T_3} = 1, m_{T_3} = 1, \ell_{T_3} = a_2$ and $\mu_{T_4} = 1, m_{T_4} = y, \ell_{T_4} = a_2$. Thus (C, T_3, T_4) is a telescoping triple.

We relabel T_3 by T_1 and T_4 by T_2 .

The -1 torus comes about from the construction. Briefly, C is obtained by performing Luttinger surgeries on the symplectic sum $(T^2 \times F_2) \#_s (T^2 \times S^2 \# 4\overline{\mathbb{CP}}^2)$ along the genus 2 surface $\{x\} \times F_2$ in $T^2 \times F_2$ and the genus 2 surface $F'_2 \subset (T^2 \times S^2) \# 4\overline{\mathbb{CP}}^2$ obtained by resolving the singularities of $(T^2 \times \{p_1\}) \cup (\{q\} \times S^2) \cup (T^2 \times \{p_2\})$ and blowing up 4 times at points on this genus 2 surface. One can choose a square zero torus of the form $T^2 \times \{y\} \subset T^2 \times F_2$ which matches up (i.e. take a relative symplectic sum) with one of the four exceptional curves to provide a -1 symplectic torus disjoint from the Lagrangian tori where the Luttinger surgeries are performed. \square

The symplectic 4-manifold $X_{1,5}$ obtained from C by $+1$ Luttinger surgeries on T_1 and T_2 as in Proposition 4 is minimal and homeomorphic to $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$ ([6]).

Our next small model is a minimal telescoping triple built in the process of constructing a minimal symplectic 4-manifold homeomorphic to $\mathbb{CP}^2 \# 7\overline{\mathbb{CP}}^2$ in [6, Theorem 8]. One stops the construction before performing the 2 Luttinger surgeries, and these unused tori provide the desired T_1 and T_2 .

Theorem 12. *There exists a minimal telescoping triple (D, T_1, T_2) with $e(D) = 10$ and $\sigma(D) = -6$. Moreover, D contains a square -1 torus disjoint from $T_1 \cup T_2$.* \square

Proof. The proof is similar to that of Theorem 11. We follow the notation and proof of [6, Theorem 8]. The 4-manifold S contains two Lagrangian tori T_1, T_2 such that $\pi_1(S - (T_1 \cup T_2))$ is generated by the two commuting elements s_1, t_1 . The Mayer-Vietoris sequence computes $H_1(S - (T_1 \cup T_2)) = \mathbb{Z}^2$ so that $\pi_1(S - (T_1 \cup T_2)) = \mathbb{Z}s_1 \oplus \mathbb{Z}t_1$.

The meridians and Lagrangian push offs of T_1 and T_2 are given by $\mu_{T_1} = 1, m_{T_1} = s_1, \ell_{T_1} = s_1^{-1}$ and $\mu_{T_2} = 1, m_{T_2} = t_1, \ell_{T_2} = s_1$. Thus (S, T_1, T_2) is a telescoping triple. It is shown to be minimal in the proof of [6, Theorem 8]. The existence of a square -1 torus follows exactly as in the proof of Theorem 11, since the manifold S is obtained by Luttinger surgeries on the symplectic sum of $(T^2 \times T^2) \# 2\overline{\mathbb{CP}}^2$ and $(T^2 \times S^2) \# 4\overline{\mathbb{CP}}^2$ along a genus 2 surface.

Relabel S as D . \square

The symplectic 4-manifold $X_{1,7}$ obtained from D by $+1$ Luttinger surgeries on T_1 and T_2 as in Proposition 4 is minimal and homeomorphic to $\mathbb{CP}^2 \# 7\overline{\mathbb{CP}}^2$ ([6]). More generally, the following proposition is true.

Proposition 13. *Let X be one of the manifolds B, B_g, C, D and T_1, T_2 the corresponding Lagrangian tori as described in Theorems 7, 11, 12, with Lagrangian push offs m_{T_i} and ℓ_{T_i} (and trivial meridians).*

Then the symplectic 4-manifolds obtained from ± 1 Luttinger surgery on one or both of T_1, T_2 along m_{T_i} or ℓ_{T_i} are all minimal.

We omit the proof, which is based on Usher's theorem and a repeated use of [6, Lemma 2]. The reader may look at the proofs of Theorems 8, 10, and 13 of [6].

Since our emphasis in this article is on 4-manifolds with odd intersection forms, we recall the following theorem due to Gompf [13].

Theorem 14. *The symplectic manifold $E'(k) = E(k)_{2,3}$ obtained from the elliptic surface $E(k)$ by performing two log transforms of order 2 and 3 is simply connected, and minimal. It has Euler characteristic $e(E'(k)) = 12k$, signature $\sigma(E'(k)) = -8k$, and an odd intersection form. \square*

6. MINIMAL SYMPLECTIC 4-MANIFOLDS WITH SIGNATURE -3 AND EULER CHARACTERISTIC GREATER THAN 14

The most complicated examples we construct are simply connected minimal symplectic 4-manifolds with signature -3 . Putting these in the context of telescoping triples is more trouble than constructing them directly. Moreover, with the exception of the $\sigma = -3$ manifolds, our inductive scheme for filling out the entire geography for $\sigma < -1$ only requires at most one copy of the manifold A of Lemma 10. Hence in this section we prove the following theorem.

Theorem 15. *For each integer $k \geq 2$, there exists a simply connected minimal symplectic 4-manifold $X_{1+2k,4+2k}$ with $e(X_{1+2k,4+2k}) = 7 + 4k$ and $\sigma(X_{1+2k,4+2k}) = -3$.*

The construction of signature -3 4-manifolds for $e = 7 + 8g$ is easier than for $e = 11 + 8g$. Roughly speaking, to produce a 4-manifold with $e = 7 + 8g$, we take the symplectic sum along a genus 3 surface of the 4-manifold A of Lemma 10 with $F \times G$, where F is a genus 3 surface and G is a genus g surface, and perform Luttinger surgery on the Lagrangian tori in G . To produce a 4-manifold with $e = 11 + 8g$ requires producing a substitute A' for A which has signature -3 and $e = 11$, and which satisfies the conclusions of Lemma 10. To do this, we take the symplectic sum of A with the product $F \times G$ of two genus 2 surfaces along a symplectic torus.

Lemma 16. *There exists a minimal symplectic 4-manifold Z with $e(Z) = 4$ and $\sigma(Z) = 0$ which contains eight homologically essential Lagrangian tori S_1, \dots, S_8 (in fact each S_i has a geometrically dual torus S_i^d so that all other intersections are zero) so that $\pi_1(Z - \cup_i S_i)$ is generated by x_1, y_1, x_2, y_2 and a_1, b_1, a_2, b_2 , and so that the meridians and Lagrangian push offs are given by*

- $S_1 : \mu_1 = [b_1^{-1}, y_1^{-1}], m_1 = x_1, \ell_1 = a_1,$
- $S_2 : \mu_2 = [x_1^{-1}, b_1], m_2 = y_1, \ell_2 = b_1 a_1 b_1^{-1},$
- $S_3 : \mu_3 = [b_2^{-1}, y_1^{-1}], m_3 = x_1, \ell_3 = a_2,$
- $S_4 : \mu_4 = [x_1^{-1}, b_2], m_4 = y_1, \ell_4 = b_2 a_2 b_2^{-1},$
- $S_5 : \mu_5 = [b_1 a_1^{-1} b_1^{-1}, y_2^{-1}], m_5 = x_2, \ell_5 = b_1^{-1},$
- $S_6 : \mu_6 = [x_2^{-1}, b_1 a_1 b_1^{-1}], m_6 = y_2, \ell_6 = b_1 a_1 b_1^{-1} a_1^{-1} b_1^{-1},$
- $S_7 : \mu_7 = [b_2 a_2^{-1} b_2^{-1}, y_2^{-1}], m_7 = x_2, \ell_7 = b_2^{-1},$
- $S_8 : \mu_8 = [x_2^{-1}, b_2 a_2 b_2^{-1}], m_8 = y_2, \ell_8 = b_2 a_2 b_2^{-1} a_2^{-1} b_2^{-1}.$

Proof. Proposition 7 of [6] (see also the construction of the manifold P in [5]) computes the fundamental group of the complement of four Lagrangian tori S_1, S_2, S_3, S_4 in the product $F_1 \times G$ of a punctured torus F_1 with a genus 2 surface G . This group is generated by loops $x_1, y_1, a_1, b_1, a_2, b_2$ (called $\tilde{x}, \tilde{y}, \tilde{a}_1, \tilde{b}_1, \tilde{a}_2, \tilde{b}_2$ there) where a_1, b_1, a_2, b_2 are a standard generating set for $\pi_1(G)$, and x_1, y_1 are a standard generating set for $\pi_1(F_1)$ based at a point h on the boundary. In particular, the boundary of $F_1 \times G$ is $\partial F_1 \times \{k\} \cup \{h\} \times G$, and the copy $\{h\} \times G$ carries the loops a_1, b_1, a_2, b_2 .

We take two copies of this manifold, calling the second $F_2 \times G$, its tori S_5, S_6, S_7, S_8 , and its generators $x_2, y_2, a'_1, b'_1, a'_2, b'_2$.

Glue the two copies together using a diffeomorphism of their boundary of the form $Id \times \phi : \partial H_2 \times G \rightarrow \partial H_1 \times G$, where $\phi : G \rightarrow G$ is the base point preserving diffeomorphism inducing the map

$$(a'_1, b'_1, a'_2, b'_2) \mapsto (b_1^{-1}, b_1 a_1 b_1^{-1}, b_2^{-1}, b_2 a_2 b_2^{-1})$$

(a composite of six Dehn twists: see [6, Lemma 9]).

The resulting manifold Z can also be described as the symplectic sum of two copies of a product of a genus 1 and genus 2 surface. Thus the result is symplectic and the 8 tori are Lagrangian. The tori S_1, S_2, S_3, S_4 in $H_1 \times G$ have geometrically dual tori $S_1^d, S_2^d, S_3^d, S_4^d$ which form a direct sum (geometrically) of four hyperbolic pairs, and similarly for S_5, S_6, S_7, S_8 . Clearly $e(Z) = 4$ and $\sigma(Z) = 0$.

Applying the Seifert-Van Kampen theorem to the formulae of Proposition 7 of [6] finishes the fundamental group assertions.

Since the diffeomorphism $Id \times \phi : \partial H_2 \times G \rightarrow \partial H_1 \times G$ extends to $H_2 \times G \rightarrow H_1 \times G$, the manifold Z is nothing but the product of two genus 2 surfaces. In particular Z is minimal. \square

Let Y be the symplectic 4-manifold obtained from the manifold Z of Lemma 16 by performing the following seven Luttinger surgeries on S_1, \dots, S_7 .

- (1) $S_1 : +1$ surgery along m_1 .
- (2) $S_2 : +1$ surgery along ℓ_2 .
- (3) $S_3 : +1$ surgery along ℓ_3 .
- (4) $S_4 : +1$ surgery along m_4 .
- (5) $S_5 : +1$ surgery along ℓ_5 .
- (6) $S_6 : +1$ surgery along m_6 .
- (7) $S_7 : +1$ surgery along m_7 .

Since the torus S_8 has not been surgered, it remains as a Lagrangian torus in Y . Since S_8 is homologically essential, the symplectic form can be perturbed so that S_8 becomes symplectic. The symplectic 4-manifold Y is minimal, since it is a symplectic sum of manifolds with contractible universal cover (see [6, Lemma 2]).

Let (B, T_1, T_2) be the telescoping triple of Theorem 7, with B containing the genus 2 symplectic surface F and geometrically dual -1 torus H_1 . Perform $+1$ Luttinger surgery on T_2 along m_{T_2} to kill t_1 , yielding a minimal (Proposition 13) symplectic 4-manifold \hat{B} . Note that \hat{B} still contains the three surfaces T_1, F, H_1 and $\pi_1(\hat{B} - (T_1 \cup F \cup H_1)) = \mathbb{Z}t_2$. The torus T_1 is disjoint from the geometrically dual symplectic surfaces F and H_1 , and its Lagrangian push offs are $m_{T_1} = 1$ and $\ell_{T_1} = t_2$, by Theorem 7.

Lemma 17. *The symplectic sum $X_{3,5} = \hat{B} \#_{T_1, S_8} Y$ is simply connected, minimal, contains a symplectic genus 2 surface of square 0 and a geometrically dual symplectic torus of square -1 . Moreover, $e(X_{3,5}) = 10$ and $\sigma(X_{3,5}) = -2$, so that $X_{3,5}$ is homeomorphic but not diffeomorphic to $3\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$.*

Proof. We refer to the notation in the statement of Theorem 7. The fundamental group of $X_{3,5} - F$ is generated by $t_2, x_1, y_1, x_2, y_2, a_1, b_1, a_2, b_2$ by the Seifert-Van Kampen theorem (recall that t_1 is killed by Luttinger surgery on T_2).

Since the meridian of T_1 in $\pi_1(B - (F \cup T_1 \cup T_2))$ is trivial, μ_8 is trivial in $\pi_1(X_{3,5} - F)$. Choose the gluing map $S_8 \rightarrow T_1$ so that ℓ_8 is killed and m_8 is sent to t_2 (i.e. $m_8 \mapsto \ell_{T_1}, \ell_8 \mapsto m_{T_1}^{-1}$).

Since ℓ_8 is a conjugate of b_2^{-1} and $m_8 = y_2$, it follows that $b_2 = 1$ and $y_2 = t_2$. This implies that μ_3 and μ_4 are trivial, and hence the third and fourth Luttinger surgeries listed above show that $a_2 = 1$ and $y_1 = 1$. Thus μ_1 and μ_7 are killed. The first and seventh Luttinger surgeries now show that $x_1 = 1$ and $x_2 = 1$. Continuing, we see that μ_2 and μ_6 are killed so that the corresponding surgeries give $a_1 = 1$ and $y_2 = 1$. This implies $\mu_5 = 1$ and so $b_1 = 1$. Hence $\pi_1(X_{3,5}) = 1$.

That $X_{3,5}$ is minimal follows from Usher's theorem. The genus 2 surface F and torus H_1 in B survive to give the required surfaces in $X_{3,5}$. \square

Proof of Theorem 15. We define two minimal simply connected symplectic 4-manifolds: let $X_- = X_{3,5}$ and let $X_+ = X_{1,3}$ (thus X_+ is obtained from the manifold \hat{B} defined above by performing +1 Luttinger surgery on T_1 along ℓ_{T_1} ; see Corollary 8). Then X_- and X_+ each contain a symplectic genus 3 surface F_3 of square 1 obtained by resolving the union $H_1 \cup F$. Moreover, $e(X_-) = 10$, $\sigma(X_-) = -2$, $e(X_+) = 6$, and $\sigma(X_+) = -2$.

Blow up X_{\pm} once at a point on F_3 and take the proper transform. Call the result \tilde{X}_{\pm} and denote by \tilde{F}_3 the proper transform of F_3 . Thus \tilde{F}_3 is a genus 3, square zero symplectic surface with simply connected complement, which meets every -1 sphere in \tilde{X}_{\pm} since X_{\pm} is minimal.

We now mimic the proof of Corollary 9. Take the product $F_3 \times G$ of a genus 3 surface with a genus g surface. Perform Luttinger surgeries on the $2g$ disjoint Lagrangian tori $Y_1 \times A_j$ and $Y_2 \times B_j$ along the curves $\ell_{Y_1 \times A_j} = a_j$ and $\ell_{Y_2 \times B_j} = b_j$ to obtain a manifold Z_g .

Then by Theorem 1 the fundamental group of Z_g is generated by the $6 + 2g$ loops $\tilde{x}_1, \tilde{y}_1, \tilde{x}_2, \tilde{y}_2, \tilde{x}_3, \tilde{y}_3, \tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_g, \tilde{b}_g$, and the relations

$$[\tilde{x}_1, \tilde{b}_j] = \tilde{a}_j, [\tilde{x}_2, \tilde{a}_j] = \tilde{b}_j$$

hold in $\pi_1(Z_g)$. Moreover, the standard symplectic generators for $\pi_1(F)$ are sent to $\tilde{x}_1, \tilde{y}_1, \tilde{x}_2, \tilde{y}_2, \tilde{x}_3, \tilde{y}_3$ in $\pi_1(Z_g)$.

Since $\pi_1(\tilde{X}_{\pm} - \tilde{F}_3) = 1$, the fundamental group of the symplectic sum

$$Q_{\pm,g} = \tilde{X}_{\pm} \#_{\tilde{F}_3, F} Z_g$$

is trivial. Indeed, the \tilde{x}_i, \tilde{y}_i are killed by taking the symplectic sum, and the relations coming from the Luttinger surgeries show the a_i and b_i are killed also.

Now $Q_{\pm,g}$ is minimal provided $g \geq 1$ by Usher's theorem since \tilde{X}_{\pm} is relatively minimal by Li's theorem, Theorem 6.

One computes:

$$e(Q_{-,g}) = e(\tilde{X}_-) + e(Z_g) + 8 = 11 + 8g - 8 + 8 = 11 + 8g,$$

$$\sigma(Q_{-,g}) = \sigma(\tilde{X}_-) + \sigma(Z_g) = -3$$

and

$$e(Q_{+,g}) = e(\tilde{X}_+) + e(Z_g) + 8 = 7 + 8g - 8 + 8 = 7 + 8g,$$

$$\sigma(Q_{+,g}) = \sigma(\tilde{X}_+) + \sigma(Z_g) = -3$$

Thus we set $X_{1+2k,4+2k} = Q_{+,k/2}$ if k is even and $X_{1+2k,4+2k} = Q_{-, (k-1)/2}$ if k is odd.

This completes the proof of Theorem 15. \square

Remark. In the construction of the manifold $X_{5,8}$, the first step (see the paragraph preceding Lemma 17) involves Luttinger surgery on the torus T_2 to kill t_1 . If one constructs the manifold $P_{5,8}$ by the same construction as for $X_{5,8}$ except by *not* performing this surgery, then $P_{5,8}$ is a minimal symplectic 4-manifold with $\pi_1(P_{5,8}) = \mathbb{Z}t_1$ containing an essential Lagrangian (or if desired symplectic) torus $T = T_2$ such that the inclusion map $\pi_1(T) \rightarrow \pi_1(P_{5,8})$ is a surjection and the inclusion map $\pi_1(P_{5,8} - T) \rightarrow \pi_1(P_{5,8})$ an isomorphism. Moreover $e(P_{5,8}) = 15$ and $\sigma(P_{5,8}) = -3$.

More generally, for any $k \geq 2$ the same construction yields a minimal symplectic 4-manifold $P_{1+2k,4+2k}$ containing a Lagrangian or symplectic torus T with these properties and such that $e(P_{1+2k,4+2k}) = 7 + 4k$, $\sigma(P_{1+2k,4+2k}) = -3$.

7. SMALL EXAMPLES WITH ODD SIGNATURE

In this section, we remind the reader of some known examples of small manifolds with odd signature, and construct a few new ones.

Kotschick showed in [17] that the Barlow surface is smoothly irreducible and hence it is a minimal symplectic 4-manifold homeomorphic to $\mathbb{CP}^2 \# 8\overline{\mathbb{CP}}^2$. This manifold realizes the pair $e = 11, \sigma = -7$.

In [14], Gompf constructs small minimal symplectic 4-manifolds which contain appropriate tori. For example, the manifold Gompf calls $S_{1,1}$ is minimal, has $e = 23$ and $\sigma = -15$, and contains a symplectic torus of square zero with simply connected complement ([14, Lemma 5.5]). The minimality $S_{1,1}$ was proved by Stipsicz [29].

Gompf also constructs other minimal symplectic 4-manifolds: the manifold $R_{2,1}$ has $e = 21$ and $\sigma = -13$ and $R_{2,2}$ has $e = 19$ and $\sigma = -11$. The minimality of $R_{2,1}$ was proved by J. Park [25], and $R_{2,2}$ was proven to be minimal by Szabó [32].

In [31], Stipsicz and Szabo construct a minimal symplectic 4-manifold homeomorphic to $\mathbb{CP}^2 \# 6\overline{\mathbb{CP}}^2$, realizing $e = 9, \sigma = -5$.

In [23], the fifth author constructs a minimal simply connected symplectic 4-manifold homeomorphic to $3\mathbb{CP}^2 \# 12\overline{\mathbb{CP}}^2$, hence with $e = 17$ and $\sigma = -9$, containing a symplectic torus $T_{2,4}$ with simply connected complement. This manifold is called X_{12} in that article, we will use the notation $X_{3,12}$ here to avoid confusion.

We produce a few more small examples.

Proposition 18. *There exists a minimal simply connected symplectic 4-manifold $X_{5,10}$ homeomorphic to $5\mathbb{CP}^2 \# 10\overline{\mathbb{CP}}^2$, hence with $e = 17$ and $\sigma = -5$.*

Proof. The manifold $X_{1,3}$ of Corollary 8 contains a symplectic genus 2 surface F of square zero, and a geometrically dual symplectic torus H_1 with square -1 . Symplectically resolve $F \cup H_1$ to produce a square 1 symplectic genus 3 surface $F_3 \subset X_{1,3}$.

Blow up $X_{1,3}$ at a point on F_3 to obtain $\tilde{X}_{1,3}$ and take \tilde{F}_3 to be the proper transform of F_3 . Then \tilde{F}_3 is a square zero symplectic surface that meets every -1 sphere in $\tilde{X}_{1,3}$ by Li's theorem (Theorem 6). Moreover, since $X_{1,3}$ is simply connected and \tilde{F}_3 meets the exceptional sphere, $\tilde{X}_{1,3} - \tilde{F}_3$ is simply connected.

Take $Y = T \times F_2$, the product of a torus with a genus 2 surface. Then Y contains the geometrically dual symplectic surfaces $T \times \{p\}$ and $\{q\} \times F_2$. Symplectically resolve their union to obtain a genus 3, square 2 symplectic surface $F'_3 \subset Y$. Note that the homomorphism induced by inclusion $\pi_1(F'_3) \rightarrow \pi_1(Y)$ is surjective. Blow up Y twice at points on F'_3 to obtain \tilde{Y} and the proper transform \tilde{F}'_3 , a square zero genus 3 symplectic surface.

Then the symplectic sum

$$X_{5,10} = \tilde{X}_{1,3} \#_{\tilde{F}_3, \tilde{F}'_3} \tilde{Y}$$

is simply connected. It is minimal by Usher's theorem.

Its characteristic classes are computed

$$e(X_{5,10}) = e(\tilde{X}_{1,3}) + e(\tilde{Y}) + 8 = 7 + 2 + 8 = 17$$

and

$$\sigma(X_{5,10}) = \sigma(\tilde{X}_{1,3}) + \sigma(\tilde{Y}) = -3 - 2 = -5.$$

The proposition follows. \square

Proposition 19. *There exists a minimal simply connected symplectic 4-manifold $X_{5,12}$ homeomorphic to $5\mathbb{CP}^2 \# 12\overline{\mathbb{CP}}^2$, hence with $e = 19$ and $\sigma = -7$.*

Proof. The proof is very similar to the proof of Proposition 18. Construct $\tilde{X}_{1,3}$ and \tilde{F}_3 as in that proof.

Take $Z = T \times T$, the product of two tori. Pick three distinct points p_1, p_2, q in T . Then Z contains the three symplectic surfaces $T \times \{p_1\}$, $T \times \{p_2\}$ and $\{q\} \times T$. Symplectically resolve their union to obtain a genus 3, square 4 symplectic surface $F'_3 \subset Z$. Note that the homomorphism induced by inclusion $\pi_1(F'_3) \rightarrow \pi_1(Z)$ is surjective. Blow up Z four times at points on F'_3 to obtain \tilde{Z} and the proper transform \tilde{F}'_3 , a square zero genus 3 symplectic surface.

Then the symplectic sum

$$X_{5,12} = \tilde{X}_{1,3} \#_{\tilde{F}_3, \tilde{F}'_3} \tilde{Z}$$

is simply connected. It is minimal by Usher's theorem.

Its characteristic classes are computed

$$e(X_{5,12}) = e(\tilde{X}_{1,3}) + e(\tilde{Z}) + 8 = 7 + 4 + 8 = 19$$

and

$$\sigma(X_{5,12}) = \sigma(\tilde{X}_{1,3}) + \sigma(\tilde{Z}) = -3 - 4 = -7.$$

The proposition follows. □

Proposition 20. *There exists a minimal simply connected symplectic 4-manifold $X_{5,14}$ homeomorphic to $5\mathbb{CP}^2 \# 14\overline{\mathbb{CP}}^2$, hence with $e = 21$ and $\sigma = -9$.*

Proof. The proof is very similar to the proof of Proposition 19. Construct $\tilde{X}_{1,3}$ and \tilde{F}_3 as in that proof.

Take $Z = T \times S^2$, the product of a torus and a sphere. Pick three distinct points p_1, p_2, p_3 in S^2 and $q \in T$. Then Z contains the four symplectic surfaces $T \times \{p_1\}, T \times \{p_2\}, T \times \{p_3\}$ and $\{q\} \times S^2$. Symplectically resolve their union to obtain a genus 3, square 6 symplectic surface $F'_3 \subset Z$. Note that the homomorphism induced by inclusion $\pi_1(F'_3) \rightarrow \pi_1(Z)$ is surjective. Blow up Z six times at points on F'_3 to obtain \tilde{Z} and the proper transform \tilde{F}'_3 , a square zero genus 3 symplectic surface.

Then the symplectic sum

$$X_{5,14} = \tilde{X}_{1,3} \#_{\tilde{F}_3, \tilde{F}'_3} \tilde{Z}$$

is simply connected. It is minimal by Usher's theorem.

Its characteristic classes are computed

$$e(X_{5,14}) = e(\tilde{X}_{1,3}) + e(\tilde{Z}) + 8 = 7 + 6 + 8 = 21$$

and

$$\sigma(X_{5,14}) = \sigma(\tilde{X}_{1,3}) + \sigma(\tilde{Z}) = -3 - 6 = -9.$$

The proposition follows. □

8. THE MAIN THEOREM

In this section we prove the first theorem stated in the introduction. We begin with an arithmetic lemma. The purpose of this lemma is to produce the number of each of the model manifolds $B, B_g, C, D, E(k)$ needed to construct 4-manifolds with specified signature and Euler characteristic. The proof includes an algorithm for finding these numbers.

Lemma 21. *Given any pair of non-negative integers (m, n) such that*

$$0 \leq m \leq 4n - 1$$

there exist non-negative integers b, c, d, g , and k so that

$$m = d + 2c + 3b + 4g \text{ and } n = b + c + d + k + g$$

and so that $b \geq 1$ if $g > 0$.

Proof. If $m = 0$, set $k = n$ and $b = c = d = g = 0$.

Assume then that $m > 0$. Choose a non-negative integer ℓ so that $\frac{m+1}{4} \leq n - \ell \leq m$. Let

$$s = \max\{z \in \mathbb{Z} \mid 3z \leq 4n - 4\ell - m - 1\} \quad \text{and} \quad \Delta = 4n - 4\ell - m - 1 - 3s.$$

Then $\Delta = 0, 1$ or 2 , and $s \geq 0$. Moreover

$$n - \ell - 1 - s = \frac{1}{3}(m - (n - \ell)) - \frac{2}{3} + \frac{\Delta}{3} \geq \frac{\Delta}{3} - \frac{2}{3}.$$

If $\Delta = 0$, then set $b = 1, c = 0, d = s, g = n - \ell - s - 1$, and $k = \ell$. Since $\Delta = 0$, and since g is an integer, $g \geq 0$.

If $\Delta = 1$ and $s \geq 1$, then set $b = 1, c = 0, d = s - 1, g = n - \ell - s - 1$, and $k = \ell + 1$. Note that $g \geq -\frac{1}{3}$ so that $g \geq 0$.

If $\Delta = 1$ and $s = 0$, then either $n - \ell - 2 \geq 0$ in which case we set $b = 2, c = 0, d = 0, k = \ell$, and $g = n - \ell - 2$, or else $n - \ell - 2 = -1$ in which case we take $b = 0, c = 1, d = 0, k = \ell$, and $g = 0$.

If $\Delta = 2$ and $s \geq 2$, set $b = 1, c = 0, d = s - 2, g = n - \ell - s - 1$ and $k = \ell + 2$.

If $\Delta = 2, s = 1$, and $n - \ell \geq 3$ then set $b = 2, c = 0, d = 0, g = n - \ell - 3$ and $k = \ell + 1$. If $\Delta = 2, s = 1$, and $n - \ell < 3$, then necessarily $n - \ell = 2$, and so $(m, n) = (2, \ell + 2)$ and we set $b = 0, c = 1, d = 0, k = \ell + 1, g = 0$.

This leaves the cases when $\Delta = 2$ and $s = 0$. If $n - \ell \geq 2$, set $b = 1, c = 1, d = 0, g = n - \ell - 2$, and $k = \ell$. Finally, if $n - \ell = 1$, then $(m, n) = (1, 1 + \ell)$, so we take $b = 0, c = 0, d = 1, g = 0$ and $k = \ell$. \square

We can now prove our main result. We state it in terms of $c_1^2 = 2e + 3\sigma$ and $\chi_h = \frac{1}{4}(e + \sigma)$ because it is simpler to work with these numbers than pairs (e, σ) where $e + \sigma \equiv 0 \pmod{4}$. Note that in this notation, a 4-manifold with $c_1^2 = 8\chi_h + k$ has signature k , so the line $c_1^2 = 8\chi_h - 2$ corresponds to manifolds with signature -2 .

Theorem 22. *For any pair (c, χ) of non-negative integers satisfying*

$$0 \leq c \leq 8\chi - 2$$

with the possible exceptions of $(c, \chi) = (5, 1), (9, 2), (13, 2)$, or $(13, 2)$, there exists a minimal simply connected symplectic 4-manifold $Y = X_{2\chi-1, 10\chi-c-1}$ with odd intersection form and

$$c = c_1^2(Y) \quad \text{and} \quad \chi = \chi_h(Y).$$

Hence Y is homeomorphic but not diffeomorphic to $(2\chi - 1)\mathbb{CP}^2 \# (10\chi - c - 1)\overline{\mathbb{CP}}^2$.

Proof. We make extensive use of the manifolds $A, B, B_g, C, D, E'(k)$ of (respectively) Lemma 10, Theorem 7, Corollary 9, Theorem 11, Theorem 12, and Theorem 14. We will also use the sporadic examples of Section 7.

We first realize all pairs with c even. Let $(m, n) = (\frac{1}{2}c, \chi)$. Lemma 21 produces integers b, c, d, g , and k so that $m = d + 2c + 3b + 4g$ and $n = b + c + d + k + g$ and so that $b \geq 1$ if $g > 0$.

Construct the symplectic sum along tori of

- (1) b copies of B if $g = 0$, or one copy of B_g and $b - 1$ copies of B if $g \geq 1$,
- (2) c copies of C , and
- (3) d copies of D .

More precisely, each of the manifolds B, C, D contain two essential Lagrangian tori. Construct the symplectic sum Z of these manifolds by chaining them together, using Proposition 3 to ensure that at each stage one has a telescoping triple.

Specifically, if $g = 0$ take

$$Z = B \#_s \cdots \#_s B \#_s C \#_s \cdots \#_s C \#_s D \#_s \cdots \#_s D$$

and if $g \geq 1$ take

$$Z = B_g \#_s B \#_s \cdots \#_s B \#_s C \#_s \cdots \#_s C \#_s D \#_s \cdots \#_s D$$

where $\#_s$ denotes the symplectic sum along the appropriate tori (perturbing the symplectic forms so that they become symplectic) according to the recipe of Theorem 3, so that the two unused Lagrangian tori (which we relabel T_1 and T_2) make (Z, T_1, T_2) a telescoping triple.

If $k = 0$, then perform +1 Luttinger surgery on T_1 and T_2 to obtain a simply connected (according to Proposition 4) symplectic 4-manifold Y .

If $k \geq 1$ and one of b, c, d is positive, perform +1 Luttinger surgery on T_2 in Z and take the symplectic sum of the result with the elliptic surface $E(k)$ along T_1 to obtain the manifold Y . Since $E(k) - T$ is simply connected, so is Y , by the same reasoning as in the proof of Proposition 4. Since B, C , and D contain +1 tori disjoint from the Lagrangian tori T_1, T_2 , the manifold Y has odd intersection form.

If $k \geq 1$ and b, c, d are zero, take $Y = E'(k)$ (see Theorem 14) which has an odd intersection form.

Thus Y is a simply connected symplectic manifold realizing the pair (c, χ) . Since each of the manifolds B, B_g, C , and D contain a surface of odd square which misses the tori used in forming the symplectic sums, and since $E'(k)$ has an odd intersection form, it follows that Y has an odd intersection form.

Since the 4-manifold Y has indefinite, odd intersection form, Freedman's theorem [12] implies that Y is homeomorphic to an appropriate connected sum of \mathbb{CP}^2 s and $\overline{\mathbb{CP}}^2$ s.

Now we turn to the case when c is odd. Suppose first that $1 \leq c \leq 8\chi - 17$. Let $(c', \chi') = (c-1, \chi-2)$. Thus $0 \leq c' \leq 8\chi' - 2$, and c' is even. Construct the manifold Z corresponding to the pair (c', χ') and either perform $+1$ Luttinger surgery on T_1 or take the symplectic sum with $E(k)$ if $k \geq 1$. But rather than performing $+1$ Luttinger surgery on T_2 as we did above, perturb the symplectic form to make T_2 symplectic, and then take the symplectic sum with Gompf's manifold $S_{1,1}$ (see Section 7) along the symplectic torus in $S_{1,1}$ with simply connected complement. Since $S_{1,1}$ has $c_1^2 = 1$ and $\chi_h = 2$ the resulting symplectic manifold Y has $(c_1^2, \chi_h) = (c, \chi)$.

Next suppose that c is odd and $7 \leq c \leq 8\chi - 11$. Set $(c', \chi') = (c-7, \chi-2)$. Thus $0 \leq c' \leq 8\chi' - 2$ and c' is even. Construct the manifold Z corresponding to the pair (c', χ') . We repeat the argument of the previous paragraph, replacing Gompf's manifold $S_{1,1}$ with the manifold $X_{3,12}$ of Section 7. Take the symplectic sum of Z with $X_{3,12}$ along T_2 and T . Since $c_1^2(X_{3,12}) = 7$ and $\chi_h(X_{3,12}) = 2$, the resulting manifold Y realizes the pair (c, χ) .

To realize all pairs (c, χ) with c odd and $21 \leq c \leq 8\chi - 5$, repeat the argument once more, this time using the manifold $P_{5,8}$ described in the remark at the end of the proof of Theorem 15, which has $c_1^2 = 21$ and $\chi_h = 3$. A bit of care must be taken to ensure that the result is simply connected since $\pi_1(P_{5,8}) = \mathbb{Z}$. This is accomplished by making sure that the generator of $\pi_1(T)$ sent to the generator of $\pi_1(P_{5,8} - T)$ is identified with an element in the kernel of $\pi_1(T_2) \rightarrow \pi_1(Z - T_2)$ when forming the symplectic sum $Y = Z \#_s P_{5,8}$.

The manifold $Y = X_{1+2k, 4+2k}$ of Theorem 15 provides an example realizing $(c, \chi) = (5 + 8k, 1 + k)$ for any $k \geq 2$, i.e. $21 \leq c = 8\chi - 3$.

Since $c_1^2 \equiv \sigma \pmod{2}$, and simply connected 4-manifolds with odd signature have an odd intersection form, it follows that the manifolds constructed for c odd also have an odd intersection form.

It remains to show that Y is minimal. Since $E'(k)$ is minimal, we assume that $c_1^2 > 0$. By Proposition 13, the 4-manifold obtained by performing one or two ± 1 Luttinger surgeries on T_1 or T_2 along ℓ_{T_i} or m_{T_i} in B, C , or D is minimal. The $E(k)$ are minimal for $k \geq 2$. Although $E(1)$ is not minimal, every -1 sphere intersects the elliptic torus. Thus Y is the symplectic sum of minimal (or, if $k = 1$, relatively minimal) symplectic 4-manifolds and therefore is minimal by Usher's theorem.

It is easy to check that the only pairs (c, χ) with $0 \leq c \leq 8\chi - 2$ which are omitted by these cases are

$$\begin{aligned} &(1, 1), (3, 1), (5, 1), \\ &(1, 2), (3, 2), (5, 2), (7, 2), (9, 2), (11, 2), (13, 2), \\ &(15, 3), (17, 3), (19, 3). \end{aligned}$$

The examples listed in Section 7 realize most of these pairs. The only ones left unrealized are $(5, 1)$, $(9, 2)$, $(13, 2)$, and $(13, 2)$.

□

The four unrealized pairs do correspond to (non-minimal) symplectic 4-manifolds; e.g. blow ups of $X_{1,3}$ or $X_{3,5}$.

It is conjectured that the irreducible smooth 4-manifold homeomorphic to $3\mathbb{CP}^2 \# 10\overline{\mathbb{CP}}^2$ constructed in [23] and the irreducible smooth 4-manifold homeomorphic to $3\mathbb{CP}^2 \# 8\overline{\mathbb{CP}}^2$ constructed in [26] are symplectic (and hence minimal): their Seiberg-Witten invariants have the right form to be the invariants of a symplectic manifold.

There exist small simply connected minimal symplectic 4-manifolds with non-negative signature (e.g. $\mathbb{CP}^2, S^2 \times S^2$). To date, no small examples are known that contain a suitable Lagrangian torus for which we can extend the construction of Theorem 22. Some moderately large examples are known and we will briefly explore the consequences for the geography problem below.

Remark. Each of the manifolds constructed in Theorem 22, with the possible exception of those corresponding $c_1^2 = 0$ and some of the small manifolds with c odd, contain nullhomologous tori suitable for altering the differentiable structure as explained in [9], using [22] to compute the change in Seiberg-Witten invariants. Those with $c_1^2 = 0$ are $E'(k)$, for which the methods of [15, 8, 11] show how to alter the differentiable structure. Hence the manifolds of Theorem 22 admit infinitely many smooth structures.

The proof of Lemma 21 provides a specific algorithm for constructing simply connected minimal 4-manifolds with desired characteristic numbers, using the model manifolds A, B, B_g, C, D , and $E(k)$.

For example, to construct a minimal symplectic manifold homeomorphic but not diffeomorphic to $3\mathbb{CP}^2 \# 17\overline{\mathbb{CP}}^2$, one sees that such a manifold would have $(c_1^2, \chi_h) = (2, 2)$. This corresponds to $(m, n) = (1, 2)$ in Lemma 21. In the notation of Lemma 21, we see that in this case $\ell = 1, s = 0$ and $\Delta = 2$, so that $b = 0, c = 0, d = 1, g = 0$, and $k = 1$. Thus the desired manifold is obtained by taking the symplectic sum

$$D \#_s E'(1)$$

and performing +1 Luttinger surgery on the remaining Lagrangian torus in D .

As another example, we construct a minimal symplectic manifold homeomorphic but not diffeomorphic to $21\mathbb{CP}^2 \# 31\overline{\mathbb{CP}}^2$, i.e. $\chi_h = 11$ and $c_1^2 = 78$. Thus $(m, n) = (39, 11)$. The proof of

Lemma 21 provides $\ell = 0$, $s = 1$ and $\Delta = 1$, and so $b = 1, c = 0, s = 1, g = 9$ and $k = 1$. Thus the desired manifold is obtained by taking the symplectic sum

$$B_9 \#_s E'(1)$$

and performing +1 Luttinger surgery on the remaining Lagrangian torus.

The integers produced by the algorithm of Lemma 21 are not unique, for example, the choice $b = 2, c = 2, d = 0, g = 8$, and $k = 0$ yields a manifold

$$B \#_s B_8 \#_s C \#_s C.$$

Performing two +1 Luttinger surgeries to this manifold yields a (possibly different) minimal symplectic manifold homeomorphic to but not diffeomorphic to $21\mathbb{CP}^2 \# 31\overline{\mathbb{CP}}^2$.

9. SIGNATURE GREATER THAN -2

Finding small minimal symplectic 4-manifolds with signature greater than -2 poses a special challenge. Stipsicz [30] shows how to produce simply connected minimal symplectic 4-manifolds with positive signature. The following theorem provides a method for producing many examples, given one. It is also useful in studying the geography problem for non-simply connected 4-manifolds.

To avoid an overly technical statement, we separate the cases of c odd and even, but a more complete statement would have the same hypotheses on (c, χ) as in Theorem 22.

Theorem 23. *Let X be a symplectic 4-manifold and suppose that X contains a symplectic torus T such that the homomorphism $\pi_1(T) \rightarrow \pi_1(X)$ induced by inclusion is trivial. Then for any pair (c, χ) of non-negative integers satisfying*

$$0 \leq c \leq 8\chi - 2 \text{ if } c \text{ is even,}$$

$$1 \leq c \leq 8\chi - 7 \text{ if } c \text{ is odd}$$

there exists a symplectic 4-manifold Y with $\pi_1(Y) = \pi_1(X)$,

$$c_1^2(Y) = c_1^2(X) + c \text{ and } \chi_h(Y) = \chi_h(X) + \chi.$$

Moreover, if X is minimal (or more generally if (X, T) is relatively minimal) then the manifold Y is minimal and has an odd, indefinite intersection form.

Proof. The argument is the same as the proof of Theorem 22 save for the last step. Let Z be as in the proof of Theorem 22. If $k = 0$, then do +1 Luttinger surgery on T_1 to get a minimal (by Proposition 13 and Usher's theorem) manifold Z_1 with $\pi_1(Z_1) \cong \mathbb{Z}$ containing a symplectic torus T_2 (after perturbing the symplectic structure) so that the induced map $\pi_1(T_2) \rightarrow \pi_1(Z_1)$ is a split surjection. If $k \geq 1$ then take a fiber sum of Z with $E(k)$ to again get a manifold Z_1

with $\pi_1(Z_1) \cong \mathbb{Z}$ containing a symplectic torus T_2 so that the induced map $\pi_1(T_2) \rightarrow \pi_1(Z_1)$ is a surjection.

Since the meridian of T_2 is nullhomotopic in Z_1 , the symplectic sum of Z_1 and X has fundamental group isomorphic to that of X , since the homomorphism $\pi_1(T) \rightarrow \pi_1(X)$ is trivial.

Minimality follows as in the proof of Theorem 22 using Usher's theorem. Since c_1^2 and χ_h are both additive with respect to symplectic sums along tori, the result follows. \square

We will require one more useful fact about B and $X_{1,3}$ not mentioned in Theorem 7 or Corollary 8, namely, the existence of a genus 2 square zero symplectic surface G geometrically dual to F . We indicate how to find G : $X_{1,3}$ is obtained by Luttinger surgery on 8 Lagrangian tori in the symplectic sum of the twice blown up 4-torus $(T^2 \times T^2) \# 2\overline{\mathbb{CP}}^2$ and the product $T \times F_2$ of a torus and a genus 2 surface.

This symplectic sum is taken along the genus 2 surface in $(T^2 \times T^2) \# 2\overline{\mathbb{CP}}^2$ obtained by resolving $T^2 \times \{p\} \cup \{q\} \times T^2$ and blowing up twice (for definiteness at points on $T^2 \times \{p\}$). In $T \times F_2$ one takes the surface $\{x\} \times F_2$.

The square -1 torus H_1 of Theorem 7 and Corollary 8 was obtained by taking the torus of the form $T \times \{z\}$ which matches up with one of the exceptional spheres in the symplectic sum. To find the surface G , take another nearby torus of the form $T \times \{z'\}$ in $T \times F_2$ and match it up with a torus of the form $\{q'\} \times T^2$. This is the required surface G . (The surface F is a parallel copy of $\{x\} \times F_2$).

Theorem 24. *For all integers $k \geq 45$, there exists a simply connected minimal symplectic 4-manifold $X_{2k+1,2k+1}$ with Euler characteristic $e = 4k + 4$ and signature $\sigma = 0$.*

For all integers $k \geq 49$ there exists a simply connected minimal symplectic 4-manifold $X_{2k-1,2k}$ with Euler characteristic $e = 4k + 1$ and signature $\sigma = -1$.

Proof. Start with the telescoping triple (B, T_1, T_2) of Theorem 7. It contains a genus 2 square zero symplectic surface F and a geometrically dual square zero symplectic genus 2 surface G . The union $F \cup G$ is disjoint from $T_1 \cup T_2$.

Perform $+1$ Luttinger surgery on T_1 along ℓ_{T_1} to kill t_2 . Call the result R . Perturb the symplectic form on R slightly so that T_2 becomes symplectic. Note that $\pi_1(R - T_2) = \pi_1(R) = \mathbb{Z}t_1, \pi_1(T_2) \rightarrow \pi_1(R)$ is surjective, and R is minimal (Proposition 13).

In [6, Theorem 18], a minimal symplectic 4-manifold $\tilde{X}_{3,5}$ homeomorphic to $3\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$ and containing a pair of symplectic tori T_3, T_4 with simply connected complement is constructed. The symplectic sum $Q = R \#_{T_2, T_3} \tilde{X}_{3,5}$ is minimal by Usher's theorem. Moreover, Q is simply connected, since $T_2 \subset R$ induces a surjection on fundamental groups. The surfaces F and G persist as square zero, symplectic geometrically dual surfaces. Since $e(Q) = 16$ and $\sigma(Q) = -4$, Q is neither rational nor ruled.

Notice that the symplectic torus T_4 in Q has simply connected complement.

In Q , take 8 parallel copies of the genus 2 surface F and one copy of G and symplectically resolve to obtain a genus 18 surface $\Sigma \subset Q$ of square 16. Blow up Q 16 times, yielding a genus 18 square zero surface $\tilde{\Sigma} \subset \tilde{Q} = Q \# 16\overline{\mathbb{CP}}^2$. By Li's theorem, every -1 sphere in \tilde{Q} intersects $\tilde{\Sigma}$. Moreover $\pi_1(\tilde{Q} - \tilde{\Sigma}) = 1$.

In [30, Lemma 2.1], a Lefschetz fibration $H \rightarrow K$ over a surface K of genus 2 is constructed which has $e = 75$ and $\sigma = 25$. This fibration admits a symplectic section of square -1 and has fiber genus 16. The 4-manifold H is an algebraic surface, and by the BMY inequality [7] is holomorphically minimal. By [16], it is also symplectically minimal. Moreover, H is neither rational nor ruled since it lies on the BMY line.

Let $\Sigma' \subset H$ denote the symplectic surface obtained by symplectically resolving the union of the fiber and section. Then Σ' has square 1, and the exact sequence of fundamental groups for a Lefschetz fibration shows that $\pi_1(\Sigma') \rightarrow \pi_1(H)$ is surjective. Blow up H once along Σ' and take the proper transform to obtain a square zero, genus 18 surface in $\tilde{\Sigma}' \subset \tilde{H} = H \# \overline{\mathbb{CP}}^2$ so that $\pi_1(\tilde{H} - \tilde{\Sigma}') \rightarrow \pi_1(\tilde{H})$ is an isomorphism and $\pi_1(\tilde{\Sigma}') \rightarrow \pi_1(\tilde{H})$ is surjective. By Li's theorem, Theorem 6, every -1 sphere in \tilde{H} intersects $\tilde{\Sigma}'$, since H is not rational or ruled.

Hence the symplectic sum $S = \tilde{Q} \#_{\tilde{\Sigma}, \tilde{\Sigma}'} \tilde{H}$ is minimal. It is simply connected since $\pi_1(\tilde{Q} - \tilde{\Sigma}) = 1$ and $\pi_1(\tilde{\Sigma}') \rightarrow \pi_1(\tilde{H})$ is surjective. Moreover, the symplectic torus $T_4 \subset S$ has simply connected complement.

Since S is the symplectic sum along a genus 18 surface,

$$e(S) = e(\tilde{Q}) + e(\tilde{H}) + 4(18 - 1) = 176$$

and

$$\sigma(S) = \sigma(\tilde{Q}) + \sigma(\tilde{H}) = 24 - 20 = 4.$$

Thus $c_1^2(S) = 364$ and $\chi_h(S) = 45$. It contains the symplectic torus T_4 with simply connected complement. Hence Theorem 23 establishes the existence of minimal, simply connected symplectic 4-manifolds

$$X_{89+2\chi, 85+10\chi-c}$$

with $c_1^2 = 364 + c$ and $\chi_h = 45 + \chi$ for any (c, χ) satisfying $0 \leq c \leq 8\chi - 2$ when c is even.

Taking $c = 8\chi - 4$ for any $\chi \geq 1$ yields $X_{89+2\chi, 89+2\chi}$, a minimal simply connected symplectic 4-manifold with signature zero. The intersection form is odd since, as one can check from Lemma 21, either $\chi = 1$ in which case the model manifold C (with its -1 torus) is used in the construction of $X_{91, 91}$, or else $\chi > 1$, in which case the model manifold B (with its -1 torus) is used in the construction of $X_{89+2\chi, 89+2\chi}$.

To get minimal symplectic 4-manifolds with signature -1 , consider the symplectic sum

$$Y = B \#_{T_1, T} P_{1+2k, 4+2k}$$

of the manifold B of Theorem 7 with the manifold $P_{1+2k,4+2k}$ of the remark at the end of Section 6 along T_1 in B and T in $P_{1+2k,4+2k}$. Since $\pi_1(T) \rightarrow \pi_1(P_{1+2k,4+2k}) = \mathbb{Z}$ is surjective, $\pi_1(P_{1+2k,4+2k} - T) \rightarrow \pi_1(P_{1+2k,4+2k})$ is an isomorphism, and $\pi_1(T_1) \rightarrow \pi_1(B)$ has image a cyclic summand, the gluing map for the symplectic sum can be chosen so that $B - \text{nb}d(T_1) \subset Y$ induces an isomorphism on fundamental groups. Hence $\pi_1(T_2) \rightarrow \pi_1(Y)$ is an isomorphism.

The symplectic sum

$$X_{93+2k,94+2k} = Y \#_{T_2, T_4} S$$

is a simply connected minimal symplectic 4-manifold with $e = 189 + 4k$ and $\sigma = -1$, for any $k \geq 2$. \square

Since any symplectic signature zero 4-manifold has e a multiple of 4, there remain 45 signature zero minimal symplectic 4-manifolds with odd intersection form to be constructed. Also missing are 48 signature -1 minimal symplectic 4-manifolds. Hence to complete the geography problem for minimal simply connected symplectic 4-manifolds of non-negative signature there remain 97 manifolds to discover.

REFERENCES

- [1] D. Auroux, S. K. Donaldson, and L. Katzarkov, *Luttinger surgery along Lagrangian tori and non-isotopy for singular symplectic plane curves*. Math. Ann. 326 (2003), no. 1, 185–203.
- [2] A. Akhmedov, *Small exotic 4-manifolds*. Preprint math.GT/0612130 (2006).
- [3] A. Akhmedov, B. Doug Park, *Exotic Smooth Structures on Small 4-Manifolds*. Preprint math.GT/0701664 (2007).
- [4] A. Akhmedov, R. I. Baykur, B. Doug Park, *Constructing infinitely many smooth structures on small 4-manifolds*. Preprint math.GT/0703480 (2007).
- [5] S. Baldridge and P. Kirk, *A symplectic manifold homeomorphic but not diffeomorphic to $\mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$* . Preprint math.GT/0702211 (2007).
- [6] S. Baldridge and P. Kirk, *Constructions of small symplectic 4-manifolds using Luttinger surgery*. Preprint math.GT/0703065 (2007).
- [7] W. Barth, C. Peters, and A. Van de Ven, “Compact complex surfaces.” Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 4. Springer-Verlag, Berlin, 1984. x+304 pp.
- [8] R. Fintushel and R. J. Stern, *Knots, links, and 4-manifolds*. Invent. Math. 134 (1998), no. 2, 363–400.
- [9] R. Fintushel, B. Doug Park, and R.J. Stern, *Reverse engineering small 4-manifolds*. Preprint, math.GT/0701846 (2007).
- [10] R. Fintushel and R. J. Stern, *Double node neighborhoods and families of simply connected 4-manifolds with $b^+ = 1$* . J. Amer. Math. Soc. 19 (2006), no. 1, 171–180.
- [11] R. Fintushel and R. J. Stern, *Surgery in cusp neighborhoods and the geography of irreducible 4-manifolds*, Invent. Math. 117 (1994), no. 3, 455–523.
- [12] M. Freedman, *The topology of four-dimensional manifolds*. J. Differential Geom. 17 (1982), no. 3, 357–453.
- [13] R. E. Gompf, *Nuclei of elliptic surfaces*. Topology 30 (1991), 479–511.
- [14] R. E. Gompf, *A new construction of symplectic manifolds*. Ann. of Math. (2) 142 (1995), no. 3, 527–595.

- [15] R. E. Gompf and T. S. Mrowka, *Irreducible 4-manifolds need not be complex*. Ann. of Math. (2) 138 (1993), no. 1, 61–111.
- [16] M. J. D. Hamilton, and D. Kotschick, *Minimality and irreducibility of symplectic four-manifolds*. Int. Math. Res. Not. 2006, Art. ID 35032, 13 pp.
- [17] D. Kotschick, *On manifolds homeomorphic to $\mathbb{CP}^2 \# 8\overline{\mathbb{CP}}^2$* . Invent. Math. 95 (1989), no. 3, 591–600.
- [18] T.-J. Li, *Smoothly embedded spheres in symplectic 4-manifolds*. Proc. Amer. Math. Soc. 127 (1999), no. 2, 609–613.
- [19] T.-J. Li and A. Liu, *Symplectic structure on ruled surfaces and a generalized adjunction formula*. Math. Res. Lett. 2 (1995), no. 4, 453–471.
- [20] K. M. Luttinger, *Lagrangian Tori in \mathbb{R}^4* . J. Diff. Geom. 52 (1999), 203–222.
- [21] D. McDuff and D. Salamon, “Introduction to symplectic topology. Second edition.” Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1998. x+486.
- [22] J. Morgan, T. Mrowka, and Z. Szabo, *Product formulas along T^3 for Seiberg-Witten invariants*. Math. Res. Lett. 4 (1997), no. 6, 915–929.
- [23] B. Doug Park, *Constructing infinitely many smooth structures on $3\mathbb{CP}^2 \# n\overline{\mathbb{CP}}^2$* . Math. Ann. 322 (2002), no. 2, 267–278, and Erratum (2007).
- [24] Jongil Park, *The geography of irreducible 4-manifolds*. Proc. Amer. Math. Soc. 126 (1998), no. 8, 2493–2503.
- [25] Jongil Park, *Exotic smooth structures on 4-manifolds*. Forum Math. 14 (2002), 915–929.
- [26] Jongil Park, *Exotic smooth structures on $3\mathbb{CP}^2 \# 8\overline{\mathbb{CP}}^2$* . Bull. London Math. Soc. 39 (2007), 95–102.
- [27] Jongil Park, *Simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$* . Invent. Math. 159 (2005), no. 3, 657–667.
- [28] Jongil Park and Ki-Heon Yun, *Exotic smooth structures on $(2n+2l-1)\mathbb{CP}^2 \# (2n+4l-1)(-\mathbb{CP}^2)$* . Preprint math.GT/0703619 (2007).
- [29] A. I. Stipsicz, *Donaldson invariants of certain symplectic manifolds*. J. Reine Angew. Math. 465 (1995), 1–10.
- [30] A. I. Stipsicz, *Simply connected symplectic 4-manifolds with positive signature*. Proceedings of 6th Gökova Geometry-Topology Conference. Turkish J. Math. 23 (1999), no. 1, 145–150.
- [31] A. I. Stipsicz and Z. Szabó, *An exotic smooth structure on $\mathbb{CP}^2 \# 6\overline{\mathbb{CP}}^2$* . Geom. Topol. 9 (2005), 813–832.
- [32] Z. Szabó, *Irreducible four-manifolds with small Euler characteristics*. Topology 35 (1996), 411–426.
- [33] C. H. Taubes, *Counting pseudo-holomorphic submanifolds in dimension 4*. J. Differential Geom. 44 (1996), no. 4, 818–893.
- [34] C. H. Taubes, *Seiberg-Witten and Gromov invariants*. Geometry and physics (Aarhus, 1995), 591–601, Lecture Notes in Pure and Appl. Math., 184, Dekker, New York, 1997.
- [35] C. H. Taubes, *The Seiberg-Witten invariants and symplectic forms*. Math. Res. Lett. 1 (1994), no. 6, 809–822.
- [36] M. Usher, *Minimality and symplectic sums*, International Mathematics Research Notices, vol. 2006, Article ID 49857, 17 pages, 2006.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY

ATLANTA, GA, 30322, USA

E-mail address: ahmadov@math.gatech.edu

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY

BATON ROUGE, LA 70817, USA

E-mail address: sbaldrid@math.lsu.edu

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY

EAST LANSING MI 48824, USA

E-mail address: baykur@msu.edu

MATHEMATICS DEPARTMENT, INDIANA UNIVERSITY

BLOOMINGTON, IN 47405, USA

E-mail address: pkirk@indiana.edu

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO

WATERLOO, ON, N2L 3G1, CANADA

E-mail address: bdpark@math.uwaterloo.ca